

MATH 152, Fall 2022
COMMON EXAM II - VERSION **B**

LAST NAME(print): _____ FIRST NAME(print): _____

INSTRUCTOR: _____

UIN: _____

SECTION NUMBER: _____

DIRECTIONS:

1. The use of a calculator, laptop or computer is prohibited.
2. TURN OFF cell phones and put them away. If a cell phone is seen during the exam, your exam will be collected and you will receive a zero.
3. In Part 1, mark the correct choice on your ScanTron using a No. 2 pencil. The scantrons will not be returned, therefore *for your own records, also record your choices on your exam!*
4. In Part 2, present your solutions in the space provided. *Show all your work* neatly and concisely and *clearly indicate your final answer*. You will be graded not merely on the final answer, but also on the quality and correctness of the work leading up to it.
5. **Be sure to fill in your name, UIN, section number and version letter of the exam on the ScanTron form.**

THE AGGIE CODE OF HONOR

“An Aggie does not lie, cheat or steal, or tolerate those who do.”

Signature: _____

Some integrals that may or may not be useful.

$$\int \sec^3 x \, dx = \frac{1}{2} \sec x \tan x + \frac{1}{2} \ln |\sec x + \tan x| + C$$

$$\text{YMLNWK} \int \csc^3 x \, dx = \frac{-1}{2} \csc x \cot x + \frac{1}{2} \ln |\csc x - \cot x| + C$$

This page left intentionally blank.

PART I: Multiple Choice. 3.5 points each

1. The sequence $a_n = \frac{(-1)^n n^2}{2n^2 + 5}$

- (a) Diverges
- (b) Converges to $\frac{1}{2}$
- (c) None of these.
- (d) Converges to 0
- (e) Converges to $-\frac{1}{2}$

2. Which of the following is an appropriate substitution to use when solving the integral $\int \sqrt{16x^2 - 9} \, dx$?

- (a) $x = \frac{3}{4} \tan \theta$
- (b) $x = \frac{3}{4} \sin \theta$
- (c) $x = \frac{4}{3} \sec \theta$
- (d) $x = \frac{4}{3} \sin \theta$
- (e) $x = \frac{3}{4} \sec \theta$

3. Which of the following is a proper Partial Fraction Decomposition for the rational function

$$\frac{5x + 1}{(x + 3)(x^2 + 4x + 3)(x^2 + 4)}$$

- (a) $\frac{A}{x + 3} + \frac{Bx + C}{x^2 + 4x + 3} + \frac{Dx + E}{x^2 + 4}$
- (b) $\frac{A}{x + 3} + \frac{Bx + C}{(x + 3)^2} + \frac{D}{x + 1} + \frac{Ex + F}{x^2 + 4}$
- (c) $\frac{A}{x + 3} + \frac{B}{(x + 3)^2} + \frac{C}{x + 1} + \frac{Dx + E}{x^2 + 4}$
- (d) $\frac{A}{x + 3} + \frac{B}{(x + 3)^2} + \frac{C}{x + 1} + \frac{D}{x + 2} + \frac{D}{x - 2}$
- (e) None of these.

4. Assume that the sequence $\{a_n\}$ is decreasing and bounded below by 1, i.e. $a_n \geq 1$, for all positive n . Determine if the sequence is convergent or divergent.

$$a_1 = 4 \quad \text{and} \quad a_{n+1} = \frac{10}{7 - a_n}$$

- (a) Convergent to 2
 - (b) Convergent to 1
 - (c) Divergent
 - (d) Convergent to $\frac{10}{7}$
 - (e) Convergent to 5
5. After an appropriate substitution, the integral $\int x^2 \sqrt{9 - x^2} dx$ is equivalent to which of the following?

(a) $81 \int \sin^2 \theta \cos^2 \theta d\theta$

(b) $81 \int \sec^3 \theta \tan^2 \theta d\theta$

(c) $27 \int \sec^2 \theta \tan \theta d\theta$

(d) $9 \int \cos^2 \theta d\theta$

(e) $27 \int \sin^2 \theta \cos \theta d\theta$

6. The series $\sum_{i=1}^{\infty} (e^{1/i} - e^{1/(i+1)})$

- (a) converges to 0
- (b) None of these.
- (c) converges to $e - 1$
- (d) diverges
- (e) converges to e

7. Let $s = \sum_{n=1}^{\infty} \frac{1}{n^3}$. Using The Remainder Estimate for the Integral Test, determine the smallest value of n that ensures that $R_n = s - s_n \leq \frac{1}{44}$.

(a) $n = 7$

(b) $n = 8$

(c) $n = 6$

(d) $n = 5$

(e) $n = 4$

8. Compute $\int_0^4 \frac{x+2}{x^2+4} dx$.

(a) $\ln 20 - \ln 4$

(b) $\frac{1}{2} (\ln 20 - \ln 4) + \arctan(2)$

(c) $\ln 6 - \ln 2$

(d) $\frac{1}{2} (\ln 20 - \ln 4) + 2 \arctan(4)$

(e) $\ln 20 - \ln 4 + 2 \arctan(4)$

9. Let $\sum_{n=1}^{\infty} a_n$ be a series whose n th partial sum is $s_n = \frac{n}{n+2}$. Find a_4 .

(a) $a_4 = \frac{2}{3}$

(b) None of these.

(c) $a_4 = \frac{1}{21}$

(d) $a_4 = \frac{1}{15}$

(e) $a_4 = 1$

10. Let $\sum_{n=1}^{\infty} a_n$ be a series whose n th partial sum is $s_n = \frac{7n^2 + 5}{5n^2 + 2}$. The series

- (a) None of these.
- (b) converges to $\frac{12}{7}$
- (c) converges to 2.5
- (d) diverges
- (e) converges to $\frac{7}{5}$

11. Which sequence is both bounded and increasing?

- (a) $a_n = \sin(2n\pi)$
- (b) $a_n = 1 - \frac{2}{n}$
- (c) $a_n = \ln n$
- (d) $a_n = e^{-n}$
- (e) None of these.

12. Compute $\int_{-1}^{\infty} \frac{1}{1+x^2} dx$.

- (a) $\frac{\pi}{4}$
- (b) None of these.
- (c) ∞
- (d) $\frac{3\pi}{4}$
- (e) $\frac{\pi}{2}$

13. The sequence $a_n = \frac{n^2}{n+2} - \frac{n^2}{n+5}$

- (a) Converges to 3
- (b) Converges to 0
- (c) None of these.
- (d) Diverges
- (e) Converges to 7

14. Compute the sum of the series $\sum_{n=1}^{\infty} \frac{(-4)^{n+1}}{5^n}$.

- (a) $\frac{16}{9}$
- (b) $\frac{-20}{9}$
- (c) None of these.
- (d) $\frac{-16}{9}$
- (e) This series diverges.

15. Which of the following statements is true regarding the improper integral $\int_1^{\infty} \frac{1}{e^x + \sqrt{x}} dx$?

- (a) The integral converges because $\int_1^{\infty} \frac{1}{e^x + \sqrt{x}} dx < \int_1^{\infty} \frac{1}{\sqrt{x}} dx$ and $\int_1^{\infty} \frac{1}{\sqrt{x}} dx$ converges.
- (b) The integral diverges because $\int_1^{\infty} \frac{1}{e^x + \sqrt{x}} dx > \int_1^{\infty} \frac{1}{e^x} dx$ and $\int_1^{\infty} \frac{1}{e^x} dx$ diverges.
- (c) The integral converges because $\int_1^{\infty} \frac{1}{e^x + \sqrt{x}} dx < \int_1^{\infty} \frac{1}{e^x} dx$ and $\int_1^{\infty} \frac{1}{e^x} dx$ converges.
- (d) The integral diverges because $\int_1^{\infty} \frac{1}{e^x + \sqrt{x}} dx > \int_1^{\infty} \frac{1}{\sqrt{x}} dx$ and $\int_1^{\infty} \frac{1}{\sqrt{x}} dx$ diverges.
- (e) The integral converges to 0.

16. Which of the following series diverges by the Test for Divergence?

(I) $\sum_{n=1}^{\infty} \cos\left(\frac{\pi n}{2n+1}\right)$

(II) $\sum_{n=1}^{\infty} \frac{4}{4 + e^{-3n}}$

(III) $\sum_{n=1}^{\infty} \frac{1}{\arctan n}$

- (a) (III) only
- (b) (II) and (III) only
- (c) (I) and (II) only
- (d) (II) only
- (e) (I), (II), and (III)

17. Which of these substitutions would be used to evaluate $\int x^2 \sqrt{x^2 + 4x + 13} dx$?

- (a) $x + 2 = 3 \sec \theta$
- (b) $x^2 + 4x = \sqrt{13} \tan \theta$
- (c) $x + 4 = \sqrt{13} \sec \theta$
- (d) none of these.
- (e) $x + 2 = 3 \tan \theta$

18. The improper integral $\int_1^e \frac{1}{x \ln x} dx$

- (a) converges to $\frac{1}{e} - 1$.
- (b) diverges to ∞ .
- (c) converges to 1.
- (d) converges to -1 .
- (e) diverges to $-\infty$.

PART II WORK OUT

Directions: Present your solutions in the space provided. *Show all your work* neatly and concisely and *Box your final answer*. You will be graded not merely on the final answer, but also on the quality and correctness of the work leading up to it.

19. (6 points) Find a general formula, a_n , for the sequence. Assume the pattern continues, and begins with $n = 1$.

$$\left\{ \frac{5}{8}, \frac{-9}{27}, \frac{13}{64}, \frac{-17}{125}, \frac{21}{216}, \dots \right\}$$

20. (5 points) Determine whether the series converges or diverges. Fully support your conclusion.

$$\sum_{n=1}^{\infty} \frac{3^{2n}}{5^n}$$

21. (6 points) Determine whether the series converges or diverges. Fully support your conclusion.

$$\sum_{n=1}^{\infty} ne^{-n^2}$$

22. (10 points) Compute $\int \frac{1}{x^4\sqrt{x^2-9}} dx$. In your final answer, any trig or inverse trig expressions that can be rewritten algebraically must be.

23. (10 points) Compute $\int \frac{x^2 + 7x + 9}{(x + 2)(x + 1)^2} dx$

DO NOT WRITE IN THIS TABLE.

Question	Points Awarded	Points
1-18		63
19		6
20		5
21		6
22		10
23		10
		100